

# A solution of a quartic equation

Tai-Choon Yoon\*

Yoon TC. A solution of a quartic equation. J Pure Appl Math. 2025;9(1): 1-3.

quartic equation as well as the solution of a quartic equation.

**Keywords:** Ferrari's solution; Resolvent cubic; Solution of a quartic equation

**ABSTRACT**

This solution is equal to L. Ferrari's if we simply change the inner square root to. This article shows the shortest way to have a resolvent cubic for a

**INTRODUCTION**

**Derivation of a solution of a quartic equation**

The solution of a quartic polynomial was discovered by Lodovico de Ferrari in 1540. Ferrari's solution is good for solving a quartic equation. This article shows a simpler way to solve the quartic equation than Ferrari [1].

A monic form of a quartic polynomial is written as

$$x^4 + c_3x^3 + c_2x^2 + c_1x + c_0 = 0. \tag{1}$$

To solve a quartic, we need to get a resolvent (1) cubic. A resolvent cubic can be obtained from the above quartic equation by using the following biquadratic equation [2].

$$(x^2 + a_1x + a_0)^2 - w(x + b_0)^2 = 0, \tag{2}$$

where  $a_1, a_0$  and  $b_0$  are arbitrary coefficients, and  $w$  is a coupling constant.

Unfolding the brackets of the equation (2) and comparing to those coefficients of the equation (1), we can find

$$\begin{aligned} a_1 &= \frac{1}{2}c_3, \\ a_0 &= \frac{1}{2}c_2 + \frac{1}{2}w - \frac{1}{8}c_3^2, \\ b_0 &= (-c_1 + \frac{1}{2}c_2c_3 + \frac{1}{2}c_3w - \frac{1}{8}c_3^3)/(2w), \end{aligned} \tag{3}$$

and we get the remaining resolvent equation

$$a_0^2 - b_0^2w - c_0 = 0. \tag{4}$$

Solving the equation (4) with substitutions from (3), we get the resolvent cubic equation with respect to  $w$ ,

$$\begin{aligned} w^3 + (2c_2 - \frac{3}{4}c_3^2)w^2 + (-4c_0 + c_1c_3 - c_2c_3^2 + c_2^2 + \frac{3}{16}c_3^4)w \\ + (c_1c_2c_3 - \frac{1}{4}c_1c_3^3 + \frac{1}{8}c_2c_3^4 - c_1^2 - \frac{1}{64}c_3^6 - \frac{1}{4}c_2^2c_3^2) = 0. \end{aligned} \tag{5}$$

As this resolvent cubic equation is somewhat lengthy and complicated, a reduced form is applicable as follows [3-5],

$$y^3 + p_1y + p_0 = 0, \tag{6}$$

where  $y$  represents

$$y = w + \frac{2}{3}c_2 - \frac{1}{4}c_3^2, \tag{7}$$

with  $p_1$  and  $p_0$  respectively

$$\begin{aligned} p_1 &= -4c_0 + c_1c_3 - \frac{1}{3}c_2^2, \\ p_0 &= \frac{8}{3}c_0c_2 - c_0c_3^2 + \frac{1}{3}c_1c_2c_3 - c_1^2 - \frac{2}{27}c_2^3. \end{aligned} \tag{8}$$

A radical solution of the cubic (6) provides

$$y = \sqrt[3]{-\frac{p_0}{2} - \sqrt{\left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3}} + \sqrt[3]{-\frac{p_0}{2} + \sqrt{\left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3}}. \tag{9}$$

Or we can get the solution in the form of  $w$  from the equation (5) by using the equation (7)

$$w = -\frac{2}{3}c_2 + \frac{1}{4}c_3^2 + \sqrt[3]{-\frac{p_0}{2} - \sqrt{\left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3}} + \sqrt[3]{-\frac{p_0}{2} + \sqrt{\left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3}}. \tag{10}$$

With  $p_1$  and  $p_0$  of (8).

It is to be noted that

$$D_4 = \left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3$$

Represents the discriminant of the above quartic equation (1) and it can be expanded as below,

Department of Applied Mathematics, Yonsei University, South Korea

**Correspondence:** Tai-Choon Yoon, Department of Applied Mathematics, Yonsei University, South Korea; E-mail: tcyoon@hanmail.net

**Received:** 15-Aug-2023, Manuscript No. PULJPAM-23-6664; **Editor assigned:** 17-Aug-2023, PreQC No. PULJPAM-23-6664 (PQ); **Reviewed:** 30-Aug-2023, QC No. PULJPAM-23-6664; **Revised:** 24-Jan-2025, Manuscript No. PULJPAM-23-6664 (R); **Published:** 31-Jan-2025, DOI: 10.37532/2752-8081.25.9(1).1-3



This open-access article is distributed under the terms of the Creative Commons Attribution Non-Commercial License (CC BY-NC) (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits reuse, distribution and reproduction of the article, provided that the original work is properly cited and the reuse is restricted to noncommercial purposes. For commercial reuse, contact [reprints@pulsus.com](mailto:reprints@pulsus.com)

$$D_4 = \left(\frac{p_0}{2}\right)^2 + \left(\frac{p_1}{3}\right)^3 \tag{11}$$

$$= -\frac{1}{108}(18c_3c_2c_1^3 - 80c_3c_2^2c_1c_0 + 144c_3^2c_2c_0^2 - 6c_3^2c_1^2c_0 - 4c_2^3c_1^2 + 16c_2^4c_0 - 192c_3c_1c_0^2 + 144c_2c_1^2c_0 - 128c_2^2c_0^2 - 27c_1^4 + 256c_0^3 + c_3^2c_2^2c_1^2 - 4c_3^2c_2^2c_0 + 18c_3^3c_2c_1c_0 - 4c_3^3c_1^3 - 27c_3^4c_0^2).$$

Now, we can derive out a radical solution of a quartic equation. It is convenient to deal with the equation (2) directly otherwise it is so complicated. It provides two quadratic solutions. One of them is [6]:

$$x^2 + (a_1 - \sqrt{w})x + a_0 - b_0\sqrt{w} = 0. \tag{12}$$

This gives two roots of the quadratic:

$$x_{1,2} = -\frac{a_1}{2} + \frac{\sqrt{w}}{2} \pm \frac{1}{2}\sqrt{(a_1 - \sqrt{w})^2 - 4(a_0 - b_0\sqrt{w})}. \tag{13}$$

Substituting with the equations (3), we get:

$$x_{1,2} = -\frac{c_3}{4} + \frac{\sqrt{w}}{2} \pm \frac{1}{4}\sqrt{3c_3^2 - 8c_2 - 4w - \frac{8c_1 - 4c_3c_2 + c_3^3}{\sqrt{w}}}. \tag{14}$$

with  $w$  from (10).

These are two roots of a quartic equation (1)

### DISCUSSION

### A full solution of a quartic equation

A general quartic equation is written as

$$c_4x^4 + c_3x^3 + c_2x^2 + c_1x + c_0 = 0. \tag{15}$$

Dividing by  $c_4$ , we get a monic quartic equation

$$x^4 + \frac{c_3}{c_4}x^3 + \frac{c_2}{c_4}x^2 + \frac{c_1}{c_4}x + \frac{c_0}{c_4} = 0. \tag{16}$$

An intermediary biquadratic equation for a solution of a general monic quartic equation (16) is given as

$$(x^2 + lx + m)^2 = w(x + n)^2. \tag{17}$$

For a reduced quartic, one may use the following form

$$(x^2 + m)^2 = w(x + n)^2,$$

this is simply equal to the above (17) in case  $l = 0$ .

Unfolding the brackets and comparing to those coefficients of the equation (17), the coefficients are given as [7]:

$$l = \frac{c_3}{2c_4}, \tag{18}$$

$$m = \frac{w}{2} - \frac{c_3^2}{8c_4^2} + \frac{c_2}{2c_4},$$

$$n = -\frac{c_3^3}{16c_4^3w} + \frac{c_3c_2}{4c_4^2w} + \frac{c_3}{4c_4} - \frac{c_1}{2c_4w},$$

Substituting these coefficients to the equation (16), we get the resolvent cubic equation,

$$x^4 + \frac{c_3}{c_4}x^3 + \frac{c_2}{c_4}x^2 + \frac{c_1}{c_4}x + R(w) = 0, \tag{19}$$

where  $R(w)$  provides the resolvent cubic with respect to  $w$ ,

$$4wR(w) = w^3 + sw^2 + tw + u = 0, \tag{20}$$

Where

$$s = -\frac{3c_3^2}{4c_4^2} + \frac{2c_2}{c_4}, \tag{21}$$

$$t = \frac{3c_3^4}{16c_4^4} - \frac{c_3^2c_2}{c_4^3} + \frac{c_3c_1}{c_4^2} + \frac{c_2^2}{c_4^2} - \frac{4c_0}{c_4},$$

$$u = -\frac{c_3^6}{64c_4^6} + \frac{c_3^4c_2}{8c_4^5} - \frac{c_3^2c_2^2}{4c_4^4} - \frac{c_3^3c_1}{4c_4^4} + \frac{c_3c_2c_1}{c_4^3} - \frac{c_1^2}{c_4^2}.$$

In case  $w=0$ , the biquadratic (17) simply becomes a perfect square of a quadratic equation  $(x^2+lx+m)^2=0$ , which includes the case  $l=0$  when it becomes  $(x^2 + m)^2=0$ .

Therefore the equation (17) is applicable for all quartic polynomials except when  $(x^2+lx+m)^2=0$  and  $(x^2+m)^2=0$ , which are simply solvable by factoring. To solve the resolvent cubic equation (20), we get a reduced form by substituting with  $w=y-s/3$ ,

$$y^3 + py + q = 0, \tag{22}$$

Where,

$$p = \frac{c_3c_1}{c_4^2} - \frac{c_2^2}{3c_4^2} - \frac{4c_0}{c_4}, \tag{23}$$

$$q = \frac{c_3c_2c_1}{3c_4^3} - \frac{c_3^2c_0}{c_4^3} - \frac{2c_2^3}{27c_4^3} + \frac{8c_2c_0}{3c_4^2} - \frac{c_1^2}{c_4^2}. \tag{24}$$

We get a solution from (22)

$$y = \sqrt[3]{-\frac{q}{2} - \sqrt{D_4}} + \sqrt[3]{-\frac{q}{2} + \sqrt{D_4}}, \tag{25}$$

where  $D_4$  is the discriminant of the quartic (15), which is given as follows,

$$D_4 = \frac{1}{4}q^2 + \frac{1}{27}p^3 \tag{26}$$

$$= -\frac{1}{108c_4^6}(18c_4c_3c_2c_1^3 - 80c_4c_3c_2^2c_1c_0 + 144c_4c_3^2c_2c_0^2 - 6c_4c_3^2c_1^2c_0 - 4c_4c_2^3c_1^2 + 16c_4c_2^4c_0 - 192c_4^2c_3c_1c_0^2 + 144c_4^2c_2c_1^2c_0 - 128c_4^2c_2^2c_0^2 - 27c_4^2c_1^4 + 256c_4^3c_0^3 + c_3^2c_2^2c_1^2 - 4c_3^2c_2^2c_0 + 18c_3^3c_2c_1c_0 - 4c_3^3c_1^3 - 27c_3^4c_0^2).$$

With these results, we have two quadratic equations that are two factors of the quartic equation (16)

$$x^2 + \left(\frac{c_3}{2c_4} - \sqrt{w}\right)x + \frac{1}{2}w + \frac{c_3^3}{16c_4^3\sqrt{w}} - \frac{c_3c_2}{4c_4^2\sqrt{w}} - \frac{c_3^2}{8c_4^2} - \frac{c_3\sqrt{w}}{4c_4} + \frac{c_2}{2c_4} + \frac{c_1}{2c_4\sqrt{w}}, \tag{27}$$

$$x^2 + \left(\frac{c_3}{2c_4} + \sqrt{w}\right)x + \frac{1}{2}w - \frac{c_3^3}{16c_4^3\sqrt{w}} + \frac{c_3c_2}{4c_4^2\sqrt{w}} - \frac{c_3^2}{8c_4^2} + \frac{c_3\sqrt{w}}{4c_4} + \frac{c_2}{2c_4} - \frac{c_1}{2c_4\sqrt{w}}. \tag{28}$$

The four roots of a quartic equation are given from the above

REFERENCES

1. Euler L. A conjecture on the forms of the roots of equations. arXiv preprint. 2008;16
2. Wikipedia contributors. Quartic equation. Wikipedia, The Free Encyclopedia; 2024.
3. Wikipedia contributors. de Moivre's formula. Wikipedia, The Free Encyclopedia; 2024.
4. Wikipedia contributors. Discriminant. Wikipedia, The Free Encyclopedia; 2024.
5. Wikipedia contributors. Trigonometric functions. Wikipedia, The Free Encyclopedia; 2024.
6. Wikipedia contributors. Hyperbolic functions. Wikipedia, The Free Encyclopedia; 2024.
7. Wikipedia contributors. Quartic. Wikipedia, The Free Encyclopedia; 2021

$$x_{1,2} = -\frac{c_3}{4c_4} + \frac{\sqrt{w}}{2} \pm \frac{1}{4c_4} \sqrt{3c_3^2 - 8c_4c_2 - 4c_4^2w - \frac{c_3^3 - 4c_4c_3c_2 + 8c_4^2c_1}{c_4\sqrt{w}}}, \quad (29)$$

$$x_{3,4} = -\frac{c_3}{4c_4} - \frac{\sqrt{w}}{2} \pm \frac{1}{4c_4} \sqrt{3c_3^2 - 8c_4c_2 - 4c_4^2w + \frac{c_3^3 - 4c_4c_3c_2 + 8c_4^2c_1}{c_4\sqrt{w}}}, \quad (30)$$

and the resolvent cubic equation of []

$$w = -\frac{2c_2}{3c_4} + \frac{c_3^2}{4c_4^2} + \frac{1}{3c_4} \sqrt[3]{c_2^3 - 36c_4c_2c_0 + \frac{27c_4c_1^2}{2} - \frac{9c_3c_2c_1}{2} + \frac{27c_3^2c_0}{2} + \frac{3\sqrt{3}}{2}\sqrt{-D_4}} + \frac{1}{3c_4} \sqrt[3]{c_2^3 - 36c_4c_2c_0 + \frac{27c_4c_1^2}{2} - \frac{9c_3c_2c_1}{2} + \frac{27c_3^2c_0}{2} - \frac{3\sqrt{3}}{2}\sqrt{-D_4}}, \quad (31)$$

with  $D_4$  of (26).

A full solution of a quartic equation is consisted of three parts of the equations (29), (31) and (26).